HOW TO COLLECT A THROAT SWAB SPECIMEN

A Handy Reference for Proper Throat Swab Specimen Collection

Have the patient tilt their head backwards, open their mouth, and stick out their tongue. Use a wooden tongue depressor to hold tongue in place.



Without touching side of the mouth use a sterile flocked swab (HydraFlock[®] #(25-3506-H) to swab the posterior nasopharynx and the tonsillar arches.

B Insert swab into sterile 1ml of Liquid Amies medium.



2



5

Break the swab handle at scored breakpoint line.

Label vial with the appropriate patient information.



There are certainly exceptions to the types of swabs and procedures described above, but we hope you'll reference this general overview whenever you need a refresher.

FOR SHIPPING AND LOGISTICS INQUIRIES, VISIT:

beckmancoulter.com/contact

FOR PRODUCT INQUIRIES, CONTACT:

+1 207-876-3130 (toll free) sales@puritanmedproducts.com

For more information, visit beckmancoulter.com/puritan

HydraFlock® is a registered trademark of Puritan Medical Products Co. LLC. ISO 13485:2016 🕻

Not all products are available in all countries.

© 2019 Beckman Coulter, Inc. All rights reserved. Beckman Coulter, the stylized logo, and the Beckman Coulter product and service marks mentioned herein are trademarks or registered trademarks of Beckman Coulter, Inc. in the United States and other countries.

For Beckman Coulter's worldwide office locations and phone numbers, please visit www.beckmancoulter.com/contact

FL-239670



